

Application of second Hankel determinant

Ali Omar

Introduction

Denote by \mathcal{A} the class of analytic functions

$$f(\xi) = \xi + a_2\xi^2 + a_3\xi^3 + \dots$$

in the open unit disk $U = \{\xi: |\xi| < 1\}$. The Hankel determinants $\mathcal{H}_j(n)$, ($n = 1, 2, 3, \dots; j = 1, 2, 3, \dots$) of f are denoted by

$$\mathcal{H}_j(n) = \begin{bmatrix} a_n & a_{n+1} & \cdot & \cdot & a_{n+j-1} \\ a_{n+1} & a_{n+2} & \cdot & \cdot & a_{n+j} \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ a_{n+j-1} & a_{n+j} & \cdot & \cdot & a_{n+2j-2} \end{bmatrix},$$

where $a_1 = 1$. Hankel determinants are beneficial, for example, in viewing that whether the certain coefficient functionals related to functions are bounded in U or not and do they carry the sharp bounds, see [1]. The applications of Hankel inequalities in the study of meromorphic functions can be seen in [2,3]. In 1966, Pommerenke [4] inspected $|\mathcal{H}_j(n)|$ of univalent functions and p -valent functions as well as starlike functions. In [5], it is evidenced that the Hankel determinants of univalent functions satisfy

$$|\mathcal{H}_j(n)| < kn^{-(1/2+\beta)j+3/2}; (n = 1, 2, 3, \dots; j = 1, 2, 3, \dots),$$

where $\beta > 0.00025$ and k depends only on j . Later, Hayman [6] demonstrated that $|\mathcal{H}_2(n)| < A_n^{1/2}$, ($n = 1, 2, 3, \dots; A$ an absolute constant) for univalent functions. Further, the Hankel determinant bounds of univalent functions with a positive Hayman index α were determined by Elhosh [7], of p -valent functions were seen in [8-10], and of close-to-convex and k -fold symmetric functions were discussed in [11]. Lately, several authors have explored the bounds $|\mathcal{H}_j(n)|$, of functions belonging to various subclasses of univalent and multivalent functions (for details, see [6, 12 – 27]). The Hankel determinant $\mathcal{H}_2(1) = a_3 - a_2^2$ is the renowned Fekete-Szegő Functional (see [28, 29]) and $\mathcal{H}_2(2)$; second, Hankel determinant is presumed by $\mathcal{H}_2(2) = a_2a_4 - a_3^2$.

An analytic function f_1 is subordinate to an analytic function f_2 , written as $f_1 < f_2$, if there is an analytic function $w: U \rightarrow U$ with $w(0) = 0$, satisfying $f_1(\xi) = f_2(w(\xi))$. Let \mathcal{P} be the class of functions with positive real part consisting of all analytic functions $p: U \rightarrow \mathbb{C}$ satisfying $p(0) = 1$ and $\text{Re}(p(z)) > 0$.

Ma and Minda [30] amalgamated various subclasses of starlike and convex functions which are subordinate to a function $\psi \in \mathcal{P}$ with $\psi(0) = 1, \psi'(0) > 0$, ψ maps U onto a region starlike with respect to 1 and symmetric with respect to real axis and familiarized the classes as below:

$$\begin{aligned}\delta^*(\psi) &= \left\{ f \in \mathcal{A}: \frac{\xi f'(\xi)}{f(\xi)} < \psi \right\} \text{ and } \mathcal{C}(\psi) \\ &= \left\{ f \in \mathcal{A}: 1 + \frac{\xi f''(\xi)}{f'(\xi)} < \psi \right\}.\end{aligned}$$

By choosing ψ satisfying Ma-Minda conditions and that maps U on to some precise regions like parabolas, cardioid, lemniscate of Bernoulli, and booth lemniscate in the righthalf of the complex plane, several fascinating subclasses of starlike and convex functions are familiarized and studied. Raina and Sokól [31] considered the class $\delta^*(\psi)$ for $\psi(z) = \xi + \sqrt{1 + \xi^2}$ and established some remarkable inequalities (also see [32] and references cited therein). Gandhi in [33] considered a class $\mathcal{S}^*(\psi)$ with $\psi = \beta e^\xi + (1 - \beta)(1 + \xi)$, $0 \leq \beta \leq 1$, a convex combination of two starlike functions. Further, coefficient inequalities of functions linked with petal type domains were widely discussed by Malik et al. ([34], see also references cited therein). The region bounded by the cardioid specified by the equation

$$(9x^2 + 9y^2 - 18x + 5)^2 - 16(9x^2 + 9y^2 - 6x + 1) = 0$$

was studied in [35]. Lately, Masih and Kanas [36] introduced novel subclasses $ST_{L(s)}$ and $CV_{L(s)}$ of starlike and convex functions, respectively. Geometrically, they consist of functions $f \in \mathcal{A}$ such that $\xi f'(\xi)/f(\xi)$ and $(\xi f'(\xi))'/f'(\xi)$, respectively, are lying in the region bounded by the limaçon

$$\begin{aligned}[(u - 1)^2 + v^2 - s^4]^2 &= 4s^2[(u - 1 + s^2)^2 + v^2] \\ \text{where } 0 < s &\leq \frac{1}{\sqrt{2}}\end{aligned}$$

Lately, Yuzaimi et al. [37] defined a region bounded by the bean-shaped limaçon region as below:

$$\begin{aligned}\Omega(U) = \{w = x + iy: (4x^2 + 4y^2 - 8x - 5)^2 \\ + 8(4x^2 + 4y^2 - 12x - 3) = 0\}, s \in [-1, 1] \setminus \{0\}.\end{aligned}$$

Suppose that

$$\varphi(\xi): U \rightarrow \mathbb{C}$$

is the function defined by

$$\varphi(\xi) = 1 + \sqrt{2}\xi + \frac{1}{2}\xi^2$$

is preferred so that the limaçon is in the bean shape [37]. Motivated by this present work and other aforesaid articles, the goal in this paper is to examine some coefficient inequalities and bounds on Hankel determinants of the Kamali-type class of starlike functions satisfying the conditions as given in Definition 1.

Definition 1. Let $\varphi: U \rightarrow \mathbb{C}$ be analytic and for $0 \leq \vartheta \leq 1$, we let the class $M(\vartheta, \varphi)$ as

$$\mathcal{M}(\vartheta, \varphi) = \left\{ f \in \mathcal{A}: \frac{9\xi^3 f'''(\xi) + (1+29)f''(\xi)\xi^2 + \xi f'(\xi)}{9\xi^2 f''(\xi) + \xi f'(\xi)} < \varphi(\xi), \xi \in U \right\},$$

where $\varphi(\xi) = 1 + \sqrt{2}\xi + (1/2)\xi^2$ as in (9).

We include the following results which are needed for the proofs of our main results.

Lemma 2 see [38]. Suppose that $p(\xi) = 1 + c_1\xi + c_2\xi^2 + \dots$, $\Re(p_1) > 0$, $\xi \in U$, then

$$\begin{aligned} |c_n| &\leq 2(n = 1, 2, 3, \dots) \\ |c_2 - vc_1^2| &\leq 2 \text{ m } \{1, |2v - 1|\} \end{aligned}$$

and the outcome is sharp for the functions formulated by

$$\begin{aligned} p(\xi) &= \frac{1 + \xi^2}{1 - \xi^2} \\ p(\xi) &= \frac{1 + \xi}{1 - \xi} \end{aligned}$$

Lemma 3 see [30]. Suppose that $p_1(\xi) = 1 + c_1\xi + c_2\xi^2 + \dots$, $\Re(p_1) > 0$, $\xi \in U$. Then,

(i) For $v < 0$ or $v > 1$, we have

$$|c_2 - vc_1^2| \leq \begin{cases} -4v + 2 & \text{if } v \leq 0 \\ 2 & \text{if } 0 \leq v \leq 1 \\ 4v - 2 & \text{if } v \geq 1 \end{cases}$$

Equality occurs when $p_1(\xi) = (1 + \xi)/(1 - \xi)$ or one of its rotations.

(ii) For $v \in (0, 1)$, the equality exists when $p_1(\xi) = (1 + \xi^2)/(1 - \xi^2)$ or one of its rotations

(iii) For $v = 0$, the equality happens when

$$p_1(\xi) = \left(\frac{1}{2} + \frac{1}{2}\vartheta\right) \frac{1 + \xi}{1 - \xi} + \left(\frac{1}{2} - \frac{1}{2}\vartheta\right) \frac{1 - \xi}{1 + \xi} \quad (0 \leq \vartheta \leq 1)$$

or one of its rotations.

Lemma 4 see [39]. If $p \in \mathcal{P}$ and is given by $p(\xi) = 1 + c_1\xi + c_2\xi^2 + \dots$ then

$$2c_2 = c_1^2 + x(4 - c^2)$$

$$4c_3 = c_1^3 + 2(4 - c_1^2)c_1x - c_1(4 - c_1^2)x^2 + 2(4 - c_1^2)(1 - |x|^2\xi)$$

for some x, ξ with $|x| \leq 1$ and $|\xi| \leq 1$.

Theorem 5. Let the function $f \in M(\vartheta, \varphi)$ be given by (1) then

$$\begin{aligned} |a_2| &\leq \frac{1}{\sqrt{2}(9 + 1)} \\ &\leq \frac{\sqrt{2}}{6(29 + 1)} \text{ m } \left\{1, \left|\frac{5}{2\sqrt{2}}\right|\right\} = \frac{5}{12(29 + 1)} \end{aligned}$$

Proof. Since $f \in M(\vartheta, \varphi)$, there exists an analytic function w with $w(0) = 0$ and $|w(\xi)| < 1$ in U such that

$$\frac{9\xi^3 f'''(\xi) + (1 + 29)f''(\xi)\xi^2 + \xi f'(\xi)}{9\xi^2 f''(\xi) + \xi f'(\xi)} = \varphi(w(\xi))$$

Define the function p_1 by

$$p_1(\xi) = \frac{1 + w(\xi)}{1 - w(\xi)} = 1 + c_1\xi + c_2\xi^2 + c_3\xi^3 + \dots$$

or, equivalently

$$\begin{aligned} w(\xi) &= \frac{p_1(\xi) - 1}{p_1(\xi) + 1} \\ &= \frac{1}{2} \left[c_1\xi + \left(c_2 - \frac{c_1^2}{2} \right) \xi^2 + \left(c_3 - c_1c_2 + \frac{c_1^3}{4} \right) \xi^3 + \dots \right] \end{aligned}$$

then p_1 is analytic in U with $p_1(0) = 1$ and has a positive real part in U . By using (20) together with (9), it is evident that

$$\begin{aligned} \varphi(w(\xi)) &= \varphi\left(\frac{p_1(\xi) - 1}{p_1(\xi) + 1}\right) = 1 + \frac{c_1\xi}{\sqrt{2}} + \left(\frac{1}{\sqrt{2}}\left(c_2 - \frac{c_1^2}{2}\right) + \frac{c_1^2}{8}\right)\xi^2 \\ &\quad + \left\{\frac{1}{\sqrt{2}}\left(c_3 - c_1c_2 + \frac{c_1^3}{4}\right) + \frac{c_1}{4}\left(c_2 - \frac{c_1^2}{2}\right)\right\}\xi^3 + \dots \end{aligned}$$

Since

$$\begin{aligned} &\frac{9\xi^3 f'''(\xi) + (1 + 2\vartheta)f''(\xi)\xi^2 + \xi f'(\xi)}{9\xi^2 f''(\xi) + \xi f'(\xi)} \\ &= 1 + 2(\vartheta + 1)a_2\xi + (-4(\vartheta + 1)^2a_2^2 + 6(2\vartheta + 1)a_3)\xi^2 \\ &\quad + [8(9 + 1)^3a_2^3 - 18(2\vartheta^2 + 3\vartheta + 1)a_2a_3 + 12(39 + 1)a_4]\xi^3 + \dots, \end{aligned}$$

and equating coefficients of ξ, ξ^2, ξ^3 from (21) to (22), we get

$$\begin{aligned} a_2 &= \frac{c_1}{2\sqrt{2}(9 + 1)}, \\ a_3 &= \frac{1}{24(2\vartheta + 1)} \left[c_1^2 \left(\frac{5}{2} - \sqrt{2} \right) + 2\sqrt{2}c_2 \right] \\ a_4 &= \frac{1}{192(3\vartheta + 1)} \left[c_1^3 \left(\frac{11}{\sqrt{2}} - 8 \right) + 4(2 - \sqrt{2})c_1c_2 + 8\sqrt{2}c_3 \right]. \end{aligned}$$

Now by applying Lemma 2, we get

$$|a_2| = \frac{1}{\sqrt{2}(\vartheta + 1)},$$

and also,

$$\begin{aligned}
|a_3| &= \frac{1}{24(2\vartheta + 1)} |2\sqrt{2}c_2 + c_1^2 \left(\frac{5}{2} - \sqrt{2}\right)| \\
&= \frac{\sqrt{2}}{12(2\vartheta + 1)} \left| c_2 - \frac{c_1^2}{2} \left(1 - \frac{5}{2\sqrt{2}}\right) \right| \\
&= \frac{\sqrt{2}}{12(2\vartheta + 1)} |c_2 - \kappa c_1^2|
\end{aligned}$$

where $\kappa = 1/2(1 - (5/2\sqrt{2}))$. Now by applying Lemma 2 , we get

$$|a_3| \leq \frac{\sqrt{2}}{6(2\vartheta + 1)} m \left\{ 1, \left| \frac{5}{2\sqrt{2}} \right| \right\} = \frac{5}{12(2\vartheta + 1)}$$

To show these bounds are sharp, we define the function $K_{\phi_n}(\xi)$, $\phi_n = q(\xi^{n-1})$ ($n = 2, 3, 4, \dots$) with $K_{\phi_n}(0) = 0 = K'_{\phi_n}(0) - 1$ by

$$\frac{9\xi^3 K'_{\phi_n}(\xi) + (1 + 29)K'_{\phi_n}(\xi)\xi^2 + \xi K'_{\phi_n}(\xi)}{9\xi^2 K'_{\phi_n}(\xi) + \xi K'_{\phi_n}(\xi)} = \varphi(\xi^{n-1})$$

Clearly, the function $K_{\phi_n} \in \mathcal{M}(\vartheta, \varphi)$. This completes the proof. Theorem 6. Let the function $f \in M(\mathcal{M}, \varphi)$ be given by (1) and for any $\omega \in \mathbb{C}$ then

$$|a_3 - \omega a_2^2| \leq \frac{\sqrt{2}}{6(2\vartheta + 1)} m \left\{ 1, \left| -\frac{5}{2\sqrt{2}} + \omega \frac{3(2\vartheta + 1)}{\sqrt{2}(\vartheta + 1)^2} \right| \right\}.$$

Proof. Let the function $f \in M(\vartheta, \varphi)$ be given by (1), as in Theorem 5, from (23) to (24), we have

$$\begin{aligned}
a_3 - \omega a_2^2 &= \frac{\sqrt{2}}{12(2\vartheta + 1)} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5}{2\sqrt{2}}\right) \right] - \omega \frac{c_1^2}{8(\vartheta + 1)^2} \\
&= \frac{\sqrt{2}}{12(2\vartheta + 1)} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5}{2\sqrt{2}} + \omega \frac{3(2\vartheta + 1)}{\sqrt{2}(\vartheta + 1)^2} \right) \right] \\
&= \frac{\sqrt{2}}{12(2\vartheta + 1)} [c_2 - \aleph c_1^2]
\end{aligned}$$

where $N = 1/2 \left(1 - (5/2\sqrt{2}) + \omega(3(2\vartheta + 1)/\sqrt{2}(\vartheta + 1)^2) \right)$.

Now by Lemma 2 , we get

$$|a_3 - \omega a_2^2| \leq \frac{\sqrt{2}}{6(2\vartheta + 1)} m \left\{ 1, \left| -\frac{5}{2\sqrt{2}} + \omega \frac{3(2\vartheta + 1)}{\sqrt{2}(\vartheta + 1)^2} \right| \right\}.$$

The result is sharp.

In particular, by taking $\omega = 1$, we get

$$|a_3 - a_2^2| \leq \frac{\sqrt{2}}{6(2\vartheta + 1)} \max \left\{ 1, \left| -\frac{5}{2\sqrt{2}} + \frac{3(29 + 1)}{\sqrt{2}(9 + 1)^2} \right| \right\}$$

Theorem 7. Let the function $f \in \mathcal{A}$ be given by (1) belongs to the class $M(\vartheta, \varphi)$ ($0 \leq \vartheta \leq 1$). Then, for any real number μ , we have

$$|a_3 - \mu a_2^2| \leq \begin{cases} \frac{5(1 + 9)^2 - 3(1 + 29)\omega}{12(29 + 1)(1 + 9)^2} & \mu \leq \delta_1 \\ \frac{\sqrt{2}}{6(29 + 1)} & \delta_1 \leq \mu \leq \delta_2 \\ \frac{3(1 + 29)\omega - 5(1 + 9)^2}{12(29 + 1)(1 + 9)^2} & \mu \geq \delta_2 \end{cases}$$

where for convenience

$$\delta_1 = \frac{(5 - 2\sqrt{2})(1 + 9)^2}{3(1 + 29)}$$

$$\delta_2 = \frac{(5 + 2\sqrt{2})(1 + 9)^2}{3(1 + 29)}$$

$$|a_3 - \mu a_2^2| + \frac{(1 + 9)^2 + (1 + 29)\mu}{1 + 2\vartheta} |a_2|^2 \leq \frac{1}{2(1 + 29)}.$$

If $\delta_3 \leq \mu \leq \delta_2$, then

$$|a_3 - \mu a_2^2| + \frac{3(1 + 9)^2 - (1 + 29)\mu}{1 + 2\vartheta} |a_2|^2 \leq \frac{1}{2(1 + 29)}.$$

These results are sharp.

Proof. Between (23) and (24) and (31), we have

$$\begin{aligned} a_3 - \omega a_2^2 &= \frac{\sqrt{2}}{12(2\vartheta + 1)} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5}{2\sqrt{2}} \right) \right] - \omega \frac{c_1^2}{8(9 + 1)^2} \\ &= \frac{\sqrt{2}}{12(2\vartheta + 1)} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5(9 + 1)^2 - 3\omega(2\vartheta + 1)}{2\sqrt{2}(\vartheta + 1)^2} \right) \right] \\ &= \frac{\sqrt{2}}{12(2\vartheta + 1)} [c_2 - \hbar c_1^2] \end{aligned}$$

where $\hbar = 1/2 \left(1 - \left((5(9 + 1)^2 - 3\omega(29 + 1))/2\sqrt{2}(9 + 1)^2 \right) \right)$. Our result now follows by virtue of Lemma 3. To show that these bounds are sharp, we define the function K_{ϕ_n} ($n = 2, 3, \dots$) by

$$\frac{9\xi^3 K'_{\phi_n}(\xi) + (1 + 2\vartheta)K'_{\phi_n}(\xi)\xi^2 + \xi K'_{\phi_n}(\xi)}{9\xi^2 K'_{\phi_n}(\xi) + \xi K'_{\phi_n}(\xi)} = \phi_n(\xi^{n-1})$$

$$K_{\phi_n}(0) = 0 = K'_{\phi_n}(0) - 1$$

and the functions F_η and G_η ($0 \leq \eta \leq 1$) by

$$\frac{9\xi^3 F'_\eta(\xi) + (1 + 2\vartheta)F'_\eta(\xi)\xi^2 + \xi F'_\eta(\xi)}{9\xi^2 F'_\eta(\xi) + \xi F'_\eta(\xi)} = \phi\left(\frac{\xi(\xi + \eta)}{1 + \eta\xi}\right) F_\eta(0)$$

$$= 0 = F'_\eta(0) - 1$$

$$\frac{9\xi^3 G'_\eta(\xi) + (1 + 2\vartheta)G'_\eta(\xi)\xi^2 + \xi G'_\eta(\xi)}{9\xi^2 G'_\eta(\xi) + \xi G'_\eta(\xi)} = \phi\left(\frac{-\xi(\xi + \eta)}{1 + \eta\xi}\right) G_\eta(0)$$

$$= 0 = G'_\eta(0) - 1.$$

Clearly, the functions $K_{\phi_n} = \varphi(\xi^{n-1})$, $F_\eta, G_\eta \in \mathcal{M}(\vartheta, \varphi)$. Also, we write $K_\phi = K_{\phi_2} = 1 + \sqrt{2}\xi + (1/2)\xi^2$. If $\mu < \delta_1$ or $\mu > \delta_2$, then the equality holds if and only if f is K_ϕ or one of its rotations. When $\delta_1 < \mu < \delta_2$, then the equality holds if and only if f is $K_{\phi_3} = \varphi(\xi^2) = 1 + \sqrt{2}\xi^2 + (1/2)\xi^4$ or one of its rotations. If $\mu = \delta_1$, then the equality holds if and only if f is F_η or one of its rotations. If $\mu = \delta_2$, then the equality holds if and only if f is G_η or one of its rotation.

2. Coefficient Estimates for the Function f^{-1}

Theorem 8. If $f \in \mathcal{M}(\vartheta, \varphi)$ and $f^{-1}(w) = w + \sum_{n=2}^{\infty} d_n w^n$ is the inverse function of f with $|w| < r_0$ where r_0 is the greater than the radius of the Koebe domain of the class $\mathcal{M}(\vartheta, \varphi)$, then for any complex number μ , we have

$$|d_2| \leq \frac{1}{\sqrt{2}(\vartheta + 1)},$$

$$|d_3| \leq \frac{\sqrt{2}}{6(2\vartheta + 1)} m \left\{ 1, \left| -\frac{5(9 + 1)^2 + 12(2\vartheta + 1)}{2\sqrt{2}(\vartheta + 1)^2} \right| \right\}.$$

Also, for any complex number μ , we have

$$|d_3 - \mu d_2^2| \leq \frac{\sqrt{2}}{6(2\vartheta + 1)} m \left\{ 1, \left| \frac{5(9 + 1)^2 + (12 + 6\mu)(2\vartheta + 1)}{2\sqrt{2}(\vartheta + 1)^2} \right| \right\}.$$

The result is sharp. In particular,

$$|d_3 - d_2^2| \leq \frac{\sqrt{2}}{6(2\vartheta + 1)} m \left\{ 1, \left| \frac{5(9 + 1)^2 + 18(2\vartheta + 1)}{2\sqrt{2}(\vartheta + 1)^2} \right| \right\}.$$

Proof. Since

$$f^{-1}(w) = w + \sum_{n=2}^{\infty} d_n w^n,$$

is the inverse function of f , we have

$$f^{-1}(f(\xi)) = f(f^{-1}(\xi)) = \xi.$$

From equations (23) to (24), we get

$$\xi + (a_2 + d_2)\xi^2 + (a_3 + 2a_2d_2 + d_3)\xi^3 + \dots = \xi$$

Equating the coefficients of ξ and ξ^2 on both sides of (45) and simplifying, we get

$$\begin{aligned} d_2 &= -a_2 = -\frac{c_1}{2\sqrt{2}(9+1)}, \\ d_3 &= 2a_2^2 - a_3 = \frac{c_1^2}{4(9+1)^2} - \frac{\sqrt{2}}{12(2\vartheta+1)} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5}{2\sqrt{2}} \right) \right] \\ &= -\frac{\sqrt{2}}{12(2\vartheta+1)} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5}{2\sqrt{2}} - \frac{6(2\vartheta+1)}{\sqrt{2}(9+1)^2} \right) \right] \\ &= -\frac{\sqrt{2}}{12(2\vartheta+1)} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5(9+1)^2 + 12(2\vartheta+1)}{2\sqrt{2}(9+1)^2} \right) \right] \end{aligned}$$

By applying Lemma 2, we get

$$\begin{aligned} |d_2| &\leq \frac{1}{\sqrt{2}(9+1)}, \\ m \left\{ 1, \left| -\frac{5(9+1)^2 + 12(2\vartheta+1)}{2\sqrt{2}(9+1)^2} \right| \right\}. \\ |d_3| &\leq \frac{\sqrt{2}}{6(2\vartheta+1)} m \left\{ 1, \left| -\frac{5(\vartheta+1)^2 + 12(2\vartheta+1)}{2\sqrt{2}(\vartheta+1)^2} \right| \right\}. \end{aligned}$$

For any complex number μ , consider

$$\begin{aligned} d_3 - \mu d_2^2 &= -\frac{\sqrt{2}}{12(2\vartheta+1)} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5(9+1)^2 + 12(2\vartheta+1)}{2\sqrt{2}(9+1)^2} \right) \right] \\ &\quad - \mu \frac{c_1^2}{8(1+9)^2} = -\frac{\sqrt{2}}{12(2\vartheta+1)} [c_2 - \rho^* c_1^2], \end{aligned}$$

where

$$\rho^* = \frac{1}{2} \left(1 - \frac{5(\vartheta+1)^2 + 3(4+\mu)(2\vartheta+1)}{2\sqrt{2}(\vartheta+1)^2} \right).$$

Taking modulus on both sides of (49) and applying Lemma 2, we get the estimate as stated in (41). This completes the proof of Theorem 8 .

3. The Logarithmic Coefficients

The logarithmic coefficients e_n of f defined in U are given by

$$\log \frac{f(\xi)}{\xi} = 2 \sum_{n=1}^{\infty} e_n \xi^n.$$

Using series expansion of $\log(1 + \xi)$ on the left hand side of (50) and equating various coefficients give

$$e_1 = \frac{a_2}{2},$$

$$e_2 = \frac{1}{2} \left(a_3 - \frac{a_2^2}{2} \right).$$

Theorem 9. Let $f \in \mathcal{M}(9, \varphi)$ with logarithmic coefficients given by (51) and (52). Then, and for any $v \in \mathbb{C}$, then

$$|e_2 - v e_1^2| \leq \frac{\sqrt{2}}{12(29+1)} \max \left\{ 1, \left| -\frac{5(9+1)^2 + (3+v)(29+1)}{2\sqrt{2}(9+1)^2} \right| \right\} \text{ get}$$

These inequalities are sharp. In particular, for $v = 1$, we

$$|e_2 - e_1^2| \leq \frac{\sqrt{2}}{12(29+1)} \max \left\{ 1, \left| -\frac{5(\vartheta+1)^2 + 4(29+1)}{2\sqrt{2}(\vartheta+1)^2} \right| \right\}$$

Proof. Using (23) and (24) in (51) and (52) and after simplification, one may have

$$e_1 = \frac{c_1}{4\sqrt{2}(\vartheta+1)}$$

$$e_2 = \frac{\sqrt{2}}{24(2\vartheta+1)} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5(\vartheta+1)^2 + 3(2\vartheta+1)}{2\sqrt{2}(\vartheta+1)^2} \right) \right]. \text{ form}$$

To determine the bounds for e_2 , we express (57) in the

$$e_2 = \frac{\sqrt{2}}{24(2\vartheta+1)} [c_2 - \mu^* c_1^2]$$

where

$$\mu^* = \left(1 - \frac{5(\vartheta+1)^2 + 3(2\vartheta+1)}{2\sqrt{2}(\vartheta+1)^2} \right)$$

then by applying Lemma 2, we get

$$|e_2| \leq \frac{\sqrt{2}}{12(2\vartheta + 1)} \max \left\{ 1, \left| \frac{5(\vartheta + 1)^2 + 3(2\vartheta + 1)}{2\sqrt{2}(\vartheta + 1)^2} \right| \right\}$$

For any $v^* \in \mathbb{C}$, from (56) to (57), we have

$$\begin{aligned} e_2 - ve_1^2 &= \frac{\sqrt{2}}{24(2\vartheta + 1)} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5(\vartheta + 1)^2 + 3(2\vartheta + 1)}{2\sqrt{2}(\vartheta + 1)^2} \right) \right] \\ &\quad - v \frac{c_1^2}{32(\vartheta + 1)^2} = \frac{\sqrt{2}}{24(2\vartheta + 1)} \\ &\quad \cdot \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5(\vartheta + 1)^2 + (3 + v)(2\vartheta + 1)}{2\sqrt{2}(\vartheta + 1)^2} \right) \right] \\ &= \frac{\sqrt{2}}{24(2\vartheta + 1)} [c_2 - \mu_1^* c_1^2] \end{aligned}$$

where

$$\mu_1^* = \frac{1}{2} \left(1 - \frac{5(\vartheta + 1)^2 + (3 + v)(2\vartheta + 1)}{2\sqrt{2}(\vartheta + 1)^2} \right).$$

An application of Lemma 2 gives the desired estimate.

4. Coefficients Associated with $\xi/f(\xi)$

In this section, we determine the coefficient bounds and Fekete-Szegő problem associated with the function $H(\xi)$ given by

$$H(\xi) = \frac{\xi}{f(\xi)} = 1 + \sum_{n=1}^{\infty} u_n \xi^n, \xi \in U$$

where $f \in \mathbb{M}(\vartheta, \varphi)$

Theorem 10. Let $f \in M(9, \varphi)$ and $H(\xi)$ are given by (63). Then

$$|u_1| \leq \frac{1}{\sqrt{2}(9 + 1)}$$

$$|u_2| \leq \frac{\sqrt{2}}{6(29 + 1)} \max \left\{ 1, \left| -\frac{5}{2\sqrt{2}} + \frac{3(2\vartheta + 1)}{\sqrt{2}(\vartheta + 1)^2} \right| \right\}.$$

The results are sharp.

Proof. By routine calculation, one may have

$$H(\xi) = \frac{\xi}{f(\xi)} = 1 - a_2 \xi + (a_2^2 - a_3) \xi^2 + (a_2^3 + 2a_2 a_3 - a_4) \xi^3 + \dots$$

Comparing the coefficients of ξ and ξ^2 on both sides of (63) and (65), we get

$$u_1 = -a_2$$

$$u_2 = a_2^2 - a_3.$$

Using (23) and (24) in (66) and (67), we obtain

$$u_1 = -\frac{c_1}{2\sqrt{2}(9+1)}$$

By Lemma 2, we get

$$|u_1| \leq \frac{1}{\sqrt{2}(9+1)}$$

Now,

$$\begin{aligned} u_2 &= \frac{c_1^2}{8(9+1)^2} - \frac{\sqrt{2}}{12(2\vartheta+1)} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5}{2\sqrt{2}} \right) \right] \\ &= -\frac{\sqrt{2}}{12(29+1)} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5}{2\sqrt{2}} + \frac{3(2\vartheta+1)}{\sqrt{2}(\vartheta+1)^2} \right) \right] \\ &= -\frac{\sqrt{2}}{12(2\vartheta+1)} [c_2 - \aleph^* c_1^2] \end{aligned}$$

where $\aleph^* = 1/2 \left(1 - (5/2\sqrt{2}) + (3(2\vartheta+1)c_1^2/\sqrt{2}(\vartheta+1)^2) \right)$. Again by using Lemma 2, we get

$$|u_2| \leq \frac{\sqrt{2}}{6(2\vartheta+1)} \max \left\{ 1, \left| -\frac{5}{2\sqrt{2}} + \frac{3(2\vartheta+1)}{\sqrt{2}(\vartheta+1)^2} \right| \right\}$$

For any $v \in \mathbb{C}$, between (68) and (70), we get

$$\begin{aligned} u_2 - vu_1^2 &= -\frac{\sqrt{2}}{12(2\vartheta+1)} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5}{2\sqrt{2}} + \frac{3(2\vartheta+1)}{\sqrt{2}(\vartheta+1)^2} \right) \right] \\ &\quad -v \frac{c_1^2}{8(\vartheta+1)^2} = -\frac{\sqrt{2}}{12(2\vartheta+1)} \\ &\quad \cdot \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{5}{2\sqrt{2}} + \frac{3(1-v)(2\vartheta+1)}{\sqrt{2}(\vartheta+1)^2} \right) \right] \\ &\quad +v \frac{3(2\vartheta+1)c_1^2}{4\sqrt{2}(\vartheta+1)^2} = -\frac{\sqrt{2}}{12(2\vartheta+1)} [c_2 - \aleph_1^{**} c_1^2] \end{aligned}$$

That is,

$$|u_2 - vu_1^2| = \frac{1}{2(2+\lambda)} |c_2 - v^{**} c_1^2|$$

where

$$\kappa_1^* = \frac{1}{2} \left(1 - \frac{5}{2\sqrt{2}} + \frac{3(1-\nu)(29+1)}{\sqrt{2}(\vartheta+1)^2} \right).$$

The result follows by application of Lemma 2 and therefore completes the proof.

5. Second Hankel Inequality for $f \in \mathcal{M}(\vartheta, \varphi)$

Theorem 11. Let the function $f \in \mathcal{M}(\vartheta, \varphi)$ be given by (1), then

$$|a_2 a_4 - a_3^2| \leq \frac{1}{18(29+1)^2}$$

Proof. Since $f \in \mathcal{M}(\vartheta, \varphi)$, there exists an analytic function w with $w(0) = 0$ and $|w(\xi)| < 1$ in U such that,

$$\frac{9\xi^3 f'''(\xi) + (1+29)f''(\xi)\xi^2 + \xi f'(\xi)}{\vartheta \xi^2 f''(\xi) + \xi f'(\xi)} = \varphi(w(\xi))$$

Therefore, between (23), (24), and (25), we get

$$\begin{aligned} a_2 a_4 - a_3^2 &= \frac{\sqrt{2}}{768} \left[c_1^4 \left\{ (6\vartheta^2 + 49 + 1) \left(\frac{-2}{3} + \frac{2\sqrt{2}}{3} \right) \right. \right. \\ &\quad \left. \left. - \frac{1}{3\sqrt{2}} (3\vartheta^2 + 4\vartheta + 1) + (12\vartheta^2 + 4\vartheta + 1) \right. \right. \\ &\quad \left. \left. \left(\frac{-2}{3} + \frac{1}{3\sqrt{2}} \right) - \frac{2\sqrt{2}(49+1)}{3} \right\} \right. \\ &\quad \left. + \frac{2c_2 c_1^2}{3} \{ 2(12\vartheta^2 + 4\vartheta + 1) + 4(6\vartheta^2 + 4\vartheta + 1) \right. \\ &\quad \left. \left(-\sqrt{2} + \frac{1}{2} \right) \right\} + 8\sqrt{2} c_1 c_3 (2\vartheta - \\ &\quad \left. - \frac{16\sqrt{2} c_2^2}{3} (3\vartheta^2 + 49 + 1) \right]. \end{aligned}$$

By writing

$$\begin{aligned} d_1 &= \frac{8\sqrt{2}}{(\vartheta+1)(3\vartheta+1)}, \\ d_2 &= \frac{4\{(18\vartheta^2 + 8\vartheta + 2) - 2\sqrt{2}(6\vartheta^2 + 4\vartheta + 1)\}}{3(\vartheta+1)(3\vartheta+1)(2\vartheta+1)^2}, \end{aligned}$$

$$d_3 = -\frac{16\sqrt{2}}{3(29+1)^2}$$

$$\begin{aligned} d_4 = & \frac{1}{(\vartheta+1)(3\vartheta+1)(2\vartheta+1)^2} \left\{ (6\vartheta^2+4\vartheta+1) \left(\frac{-2}{3} + \frac{2\sqrt{2}}{3} \right) \right. \\ & - (3\vartheta^2+49+1) \frac{1}{3\sqrt{2}} + (12\vartheta^2+49+1) \left(\frac{-2}{3} + \frac{1}{3\sqrt{2}} \right) \\ & \left. - \frac{2\sqrt{2}(49+1)}{3} \right\} = \frac{1}{(\vartheta+1)(3\vartheta+1)(2\vartheta+1)^2} \\ & \left(\frac{28-16\sqrt{2}}{3\sqrt{2}}\vartheta^2 - \frac{11}{3}\vartheta - \frac{4}{3} \right) \end{aligned}$$

and $T = \sqrt{2}/768$, we have

$$|a_2a_4 - a_3^2| = T|d_1c_1c_3 + d_2c_1^2c_2 + d_3c_2^2 + d_4c_1^4|.$$

From (15) to (16), it follows that

$$\begin{aligned} |a_2a_4 - a_3^2| = & \frac{T}{4} |c^4(d_1 + 2d_2 + d_3 + 4d_4) + 2xc^2(4 - c^2) \\ & \cdot (d_1 + d_2 + d_3) + (4 - c^2)x^2(-d_1c^2 + d_3(4 - c^2)) \\ & + 2d_1c(4 - c^2)(1 - |x|^2)| \end{aligned}$$

Replacing $|x|$ by μ and then substituting the values of d_1, d_2, d_3 , and d_4 from (81) yield

$$\begin{aligned} |a_2a_4 - a_3^2| \leq & \frac{T}{4(\vartheta+1)(3\vartheta+1)(2\vartheta+1)^2} \\ & \cdot \left[c^4 \left((49+1) + \frac{4}{3\sqrt{2}}(12\vartheta^2+49+1) \right. \right. \\ & - \frac{4}{3\sqrt{2}}(9+1)(39+1) \left. + 16\sqrt{2}(2\vartheta+1)^2 \right. \\ & \left. c(4-c^2)(1-\mu^2) + 2\mu c^2(4-c^2) \right. \\ & \left. \left(\frac{4}{3}(12\vartheta^2+4\vartheta+1) + \frac{4}{3}(6\vartheta^2+4\vartheta+1) \right) \right. \\ & \left. - \mu^2(4-c^2) \left(\frac{8\sqrt{2}}{3}(6\vartheta^2+4\vartheta+1)c^2 \right. \right. \\ & \left. \left. + \frac{64\sqrt{2}}{3}(39+1)(9+1) \right) \right] \\ = & \frac{T}{(\vartheta+1)(3\vartheta+1)(2\vartheta+1)^2} \left[\frac{c^4}{3} - 2\sqrt{2}(4\vartheta+1) + \frac{1}{\sqrt{2}} \right] \end{aligned}$$

$$\begin{aligned}
& (12\vartheta^2 + 4\vartheta + 1) - \frac{1}{\sqrt{2}}(\vartheta + 1)(3\vartheta + 1) \mid \\
& + 4\sqrt{2}(2\vartheta + 1)^2 c(4 - c^2) - 4\sqrt{2}(2\vartheta + 1)^2 \\
& \cdot c(4 - c^2)\mu^2 + \frac{2}{3}\mu c^2(4 - c^2)(2(12\vartheta^2 + 4\vartheta + 1) \\
& + (12\vartheta^2 + 8\vartheta + 2)) - \frac{2\sqrt{2}}{3}\mu^2(4 - c^2)(c^2(6\vartheta^2 + 4\vartheta + 1) \\
& + 8(3\vartheta^2 + 4\vartheta + 2)) \Big] = \frac{T}{(\vartheta + 1)(3\vartheta + 1)(2\vartheta + 1)^2} \\
& \cdot \left[\frac{c^4}{3} \left| -2\sqrt{2}(4\vartheta + 1) + \frac{9}{\sqrt{2}}\vartheta^2 \right| + 4\sqrt{2}(2\vartheta + 1)^2 c(4 - c^2) \right. \\
& + \frac{1}{3}\mu c^2(4 - c^2)(36\vartheta^2 + 16\vartheta + 4) - \frac{2\sqrt{2}}{3}\mu^2(4 - c^2) \\
& \cdot (c^2(6\vartheta^2 + 4\vartheta + 1) + 6(2\vartheta + 1)^2 c + 8(3\vartheta^2 + 4\vartheta + 2)) \Big] \\
& \equiv F(c, \mu, \vartheta).
\end{aligned}$$

Differentiating $F(c, \mu, \vartheta)$ in (84) partially with respect to μ yields

$$\begin{aligned}
\frac{\partial F}{\partial \mu} = & \frac{T}{(\vartheta + 1)(3\vartheta + 1)(2\vartheta + 1)^2} \left[\frac{1}{3}c^2(4 - c^2)(36\vartheta^2 + 16\vartheta + 4) \right. \\
& - \frac{4\sqrt{2}}{3}\mu(4 - c^2)(c^2(6\vartheta^2 + 4\vartheta + 1) + 6(2\vartheta + 1)^2 c \\
& \left. + 8(3\vartheta^2 + 4\vartheta + 2)) \right].
\end{aligned}$$

It is clear from (85) that $\partial F / \partial \mu > 0$; thus, $F(c, \mu, \vartheta)$ is an increasing function of μ for $0 < \mu < 1$ and for any fixed c with $0 < c < 2$. So, the maximum of $F(c, \mu, \vartheta)$ occurs at $\mu = 1$ and

$$\max F(c, \mu, \vartheta) = F(c, 1, \vartheta) \equiv G(c, \vartheta).$$

Note that

Differentiating $G(c, \vartheta)$ partially with respect to c yields

$$\begin{aligned}
G'(c, \vartheta) = & \frac{T}{(\vartheta + 1)(3\vartheta + 1)(2\vartheta + 1)^2} \left[\frac{4c^3}{3} \left(\left| -2\sqrt{2}(4\vartheta + 1) + \frac{9\vartheta^2}{\sqrt{2}} \right| \right. \right. \\
& \left. - 4(9\vartheta^2 + 4\vartheta + 1) + 2\sqrt{2}(1 + 4\vartheta + 6\vartheta^2) \right) \\
& \left. + \frac{16c}{3} (3(8\vartheta^2 + 4\vartheta + 1) - \sqrt{2}(12\vartheta^2 + 8\vartheta + 2)) \right]
\end{aligned}$$

If $G'(c, \vartheta) = 0$ then its root is $c = 0$. Also, we have

$$G''(c, \vartheta) = \frac{T}{(\vartheta + 1)(3\vartheta + 1)(2\vartheta + 1)^2} \left[4c^2 \left(1 - 2\sqrt{2}(4\vartheta + 1) + \frac{9\vartheta^2}{\sqrt{2}} \right) - 4(9\vartheta^2 + 4\vartheta + 1 + 22\sqrt{2}(1 + 4\vartheta + 6\vartheta^2)) + \frac{16}{3}(3(8\vartheta^2 + 4\vartheta + 1) - \sqrt{2}(12\vartheta^2 + 8\vartheta + 2)) \right]$$

is negative for $c = 0$, which means that the function $G(c, \vartheta)$ can take the maximum value at $c = 0$, also which is

$$|a_2 a_4 - a_3^2| \leq G(0, 9) = \frac{1}{18(29 + 1)^2}.$$

9. References

1. M. Abramowitz and I. A. Stegun (Eds.), Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, Dover Publications Inc., New York, 1965.
2. A. A. Attiya, Some applications of Mittag-Leffler function in the unit disk, Filomat 30(7) (2016), 2075-2081. <https://doi.org/10.2298/FIL1607075A>
3. D. Bansal and J. K. Prajapat, Certain geometric properties of the Mittag-Leffler functions, Complex Var. Elliptic Equ. 61(3) (2016), 338-350. <https://doi.org/10.1080/17476933.2015.1079628>
4. T. Bulboacă and G. Murugusundaramoorthy, Univalent functions with positive coefficients involving Pascal distribution series, Commun. Korean Math. Soc. 35(3) (2020), 867 – 877. <https://doi.org/10.4134/CKMS.c190413>
5. N. E. Cho, S. Y. Woo and S. Owa, Uniform convexity properties for hypergeometric functions, Fract. Calc. Appl. Anal. 5(3) (2002), 303-313.
6. P. L. Duren, Univalent Functions, Grundlehren der Mathematischen Wissenschaften Series 259, Springer Verlag, New York, 1983. [7] M. El-Deeb, T. Bulboacă and J. Dziok, Pascal distribution series connected with certain subclasses of univalent functions, Kyungpook Math. J. **59** (2019), 301-314. <https://doi.org/10.5666/KMJ.2019.59.2.301>
7. B. A. Frasin, An application of an operator associated with generalized Mittag-Leffler function, Konuralp J. Math. 7(1) (2019), 199-202.
8. B. A. Frasin, T. Al-Hawary and F. Yousef, Some properties of a linear operator involving generalized Mittag-Leffler function, Stud. Univ. Babeş-Bolyai Math. 65(1) (2020), 67 – 75. <https://doi.org/10.24193/subbmath.2020.1.06>
9. B. A. Frasin, T. Al-Hawary and F. Yousef, Necessary and sufficient conditions for hypergeometric functions to be in a subclass of analytic functions, Afr. Mat. 30(1-2) (2019), 223-230. <https://doi.org/10.1007/s13370-018-0638-5>
10. H. J. Haubold, A. M. Mathai and R. K. Saxena, Mittag-Leffler functions and their applications, J. Appl. Math. 2011(2011), Article ID 298628. <https://doi.org/10.1155/2011/298628>
11. V. Kiryakova, Generalized Fractional Calculus and Applications, Pitman Research Notes in Mathematics Series 301, Longman Scientific & Technical, Harlow, John Wiley & Sons, Inc., New York, 1994.
12. E. Merkes and B. T. Scott, Starlike hypergeometric functions, Proc. Amer. Math. Soc. 12 (1961), 885-888.

13. G. M. Mittag-Leffler, Sur la nouvelle fonction $E(x)$, C. R. Acad. Sci. Paris 137(1903), 554 – 558.
14. G. Murugusundaramoorthy, Subordination results for spiral-like functions associated with the Srivastava-Attiya operator, Integral Transforms Spec. Funct. 23(2) (2012), 97-103. <https://doi.org/10.1080/10652469.2011.562501>
15. G. Murugusundaramoorthy, D. Răducanu and K. Vijaya, A class of spirallike functions defined by Ruscheweyh-type q -difference operator, Novi Sad J. Math. 49(2)(2019), 59 – 71. <https://doi.org/10.30755/NSJOM.08284>
16. G. Murugusundaramoorthy, K. Vijaya and S. Porwal, Some inclusion results of certain subclass of analytic functions associated with Poisson distribution series, Hacet. J. Math. Stat. 45(4) (2016), 1101-1107. <https://doi.org/10.15672/HJMS.20164513110>
17. G. Murugusundaramoorthy, Subclasses of starlike and convex functions involving Poisson distribution series, Afr. Mat. 28(2017), 1357-1366. <https://doi.org/10.1007/s13370-017-0520-x>
18. S. Porwal and M. Kumar, A unified study on starlike and convex functions associated with Poisson distribution series, Afr. Mat. 27(5) (2016), 1021-1027. <https://doi.org/10.1007/s13370-016-0398-z>
19. M. S. Robertson, On the theory of univalent functions, Ann. of Math. (2) 37(2) (1936), 374-408.
20. Salah, Jamal, Hameed Ur Rehman, and Iman Al-Buwaiqi. "The Non-Trivial Zeros of the Riemann Zeta Function through Taylor Series Expansion and Incomplete Gamma Function." *Mathematics and Statistics* 10.2 (2022): 410-418.
21. Rehman, Hameed Ur, Maslina Darus, and Jamal Salah. "Graphing Examples of Starlike and Convex Functions of order β ." *Appl. Math. Inf. Sci* 12.3 (2018): 509-515.
22. Salah, Jamal Y., and O. M. A. N. Ibra. "Properties of the Modified Caputo's Derivative Operator for certain analytic functions." *International Journal of Pure and Applied Mathematics* 109.3 (2014): 665-671.
23. Jamal Salah and Maslina Darus, On convexity of the general integral operators, An. Univ. Vest Timis. Ser. Mat. -Inform. 49(1) (2011), 117-124.
24. Salah, Jamal. "TWO NEW EQUIVALENT STATEMENTS TO RIEMANN HYPOTHESIS." (2019).
25. H. Silverman, Starlike and convexity properties for hypergeometric functions, J. Math. Anal. Appl. 172 (1993), 574-581. <https://doi.org/10.1006/jmaa.1993.1044>
26. H. M. Srivastava, G. Murugusundaramoorthy and S. Sivasubramanian, Hypergeometric functions in the parabolic starlike and uniformly convex domains, Integral Transforms Spec. Funct. 18 (2007), 511 – 520. <https://doi.org/10.1080/10652460701391324>
27. L. Späček, Contribution à la théorie des fonctions univalentes, Časopis Pro Pěstování Matematiky 62(1932), 12 – 19.
28. Salah, Jamal Y. "A new subclass of univalent functions defined by the means of Jamal operator." *Far East Journal of Mathematical Sciences (FJMS)* Vol 108 (2018): 389-399.
29. Jamal Y. Salah On Riemann Hypothesis and Robin's Inequality. International Journal of Scientific and Innovative Mathematical Research (IJSIMR). Volume (3) 4 (2015) 9-14.
30. Salah, Jamal. "Neighborhood of a certain family of multivalent functions with negative coefficients." *Int. J. Pure Appl. Math* 92.4 (2014): 591-597.
31. Salah, Jamal, and Maslina Darus. "A note on Starlike functions of order α associated with a fractional calculus operator involving Caputo's fractional." *J. Appl. comp Sc. Math* 5.1 (2011): 97-101.

32. J. Salah, Certain subclass of analytic functions associated with fractional calculus operator, Trans. J. Math. Mech., 3 (2011), 35–42. Available from: <http://tjmm.edyopress.ro/journal/11030106.pdf>.
33. Salah, Jamal. "Some Remarks and Propositions on Riemann Hypothesis." *Mathematics and Statistics* 9.2 (2021): 159-165.
34. Salah, Jamal Y. Mohammad. "The consequence of the analytic continuity of Zeta function subject to an additional term and a justification of the location of the non-trivial zeros." *International Journal of Science and Research (IJSR)* 9.3 (2020): 1566-1569.
- 35.
36. A. Swaminathan, Certain sufficient conditions on Gaussian hypergeometric functions, Journal of Inequalities in Pure and Applied Mathematics 5(4) (2004), Article ID 83, 10 pages.
37. A. Wiman, Über die nullstellen der funktionen $E_\alpha(x)$, Acta Math. 29 (1905), 217-134.
38. Salah, Jamal Y. Mohammad. "An Alternative perspective to Riemann Hypothesis." *PSYCHOLOGY AND EDUCATION* 57.9 (2020): 1278-1281.
39. Salah, Jamal Y. "A note on the Hurwitz zeta function." *Far East Journal of Mathematical Sciences (FJMS)* 101.12 (2017): 2677-2683.
40. Jamal Y. Salah. Closed-to-Convex Criterion Associated to the Modified Caputo's fractional Calculus Derivative Operator. Far East Journal of Mathematical Sciences (FJMS). Vol. 101, No. 1, pp. 55-59, 2017, DOI: 10.17654/MS101010055.
41. Jamal Y. Salah, A Note on Riemann Zeta Function, IOSR Journal of Engineering (IOSRJEN), vol. 06, no. 02, pp. 07-16, February 2016, URL: [http://iosrjen.org/Papers/vol6_issue2%20\(part-3\)/B06230716.pdf](http://iosrjen.org/Papers/vol6_issue2%20(part-3)/B06230716.pdf)
42. Jamal Y. Salah, A Note on Gamma Function, International Journal of Modern Sciences and Engineering Technology (IJMSET), vol. 2, no. 8, pp. 58-64, 2015.
43. Salah, Jamal, and S. Venkatesh. "Inequalities on the Theory of Univalent Functions." *Journal of Mathematics and System Science* 4.7 (2014).
44. T. R. Prabhakar, A single integral equation with a generalized Mittag – Leffler function in the kernel, Yokohama Math. J. 19 (1971), 7 – 15.
45. [Omar Jawabreh](#), [Ahmad Abdel Qader](#), [Jamal Salah](#), [Khaled Al Mashrafi](#), [Emad Al Dein](#) [AL Fahmawee](#), [Basel J. A. Ali](#), Fractional Calculus Analysis of Tourism Mathematical Model, Progr. Fract. Differ. Appl. Vol. 9, No. S1, pp. 1 -11 202) DOI: 10.18576/pfda/09S101 URL:
46. Jamal Salah. Subordination and superordination involving certain fractional operator. Asian Journal of Fuzzy and Applied Mathematics, vol. 1, pp. 98-107, 2013. URL: <https://www.ajouronline.com/index.php/AJFAM/article/view/724>
47. Salah, Jamal Y. Mohammad. "Two Conditional proofs of Riemann Hypothesis." *International Journal of Sciences: Basic and Applied Research (IJSBAR)* 49.1 (2020): 74-83.
48. Salah, Jamal. "Fekete-Szego problems involving certain integral operator." *International Journal of Mathematics Trends and Technology-IJMTT* 7 (2014).

49. K. I. Noor, "Hankel determinant problem for the class of functions with bounded boundary rotation," *Revue Roumaine de Mathématiques Pures et Appliquées*, vol. 28, no. 8, pp. 731-739, 1983.
50. K. I. Noor and I. M. A. Al-Naggar, "On the Hankel determinant problem," *Journal of Natural Geometry*, vol. 14, no. 2, pp. 133-140, 1998.
51. R. M. Ali, S. K. Lee, V. Ravichandran, and S. Supramaniam, "The Fekete-Szegő coefficient functional for transforms of analytic functions," *Bulletin of the Iranian Mathematical Society*, vol. 35, no. 2, pp. 119-142, 2009.
52. R. M. Ali, V. Ravichandran, and N. Seenivasagan, "Coefficient bounds for p -valent functions," *Applied Mathematics and Computation*, vol. 187, no. 1, pp. 35-46, 2007.
53. W. C. Ma and D. Minda, "A united treatment of some special classes of univalent functions," in *Proceedings of the Conference on Complex Analysis*, Tianjin, 1992.
54. R. K. Raina and J. Sokół, "On coefficient estimates for a certain class of starlike functions, Hacet," *Journal of Mathematics and Statistics*, vol. 44, no. 6, pp. 1427-1433, 2015.
55. G. Murugusundaramoorthy and T. Bulboacă, "Hankel determinants for new subclasses of analytic functions related to a shell shaped region," *Mathematics*, vol. 8, no. 6, article 1041, 2020.
56. S. Gandhi, "Radius Estimates for Three Leaf Function and Convex Combination of Starlike Functions," in *Mathematical Analysis I: Approximation Theory. ICRAPAM 2018*, vol. 306 of *Springer Proceedings in Mathematics & Statistics*, Springer, Singapore.
57. S. N. Malik, S. Mahmood, M. Raza, S. Farman, and S. Zainab, "Coefficient inequalities of functions associated with petal type domains," *Mathematics*, vol. 6, no. 12, p. 298, 2018.
58. K. Sharma, N. K. Jain, and V. Ravichandran, "Starlike functions associated with a cardioid," *Afrika Matematika*, vol. 27, pp. 923-939, 2016.
59. V. S. Masih and S. Kanas, "Subclasses of starlike and convex functions associated with the Limaçon domain," *Symmetry*, vol. 12, no. 6, p. 942, 2020.
60. Y. Yuzaimi, A. H. Suzeini, and A. B. Akbarally, "Subclass of starlike functions associated with a limaçon," *AIP Conference Proceedings*, vol. 1974, article 030023, 2018.
61. P. L. Duren, *Univalent Functions. Grundlehren der Mathematischen Wissenschaften*, vol. 259, Springer, New York, 1983.
62. U. Grenander and G. Szegő, *Toeplitz Forms and Their Applications. California Monographs in Mathematical Sciences*, University of California Press, Berkeley, 1958.